

## Sawtail Shark, *Figaro boardmani*

Report Card assessment	Sustainable		
IUCN Red List Australian Assessment	Endemic to Australia	IUCN Red List Global Assessment	Least Concern
Assessors	Kyne, P.M. & Bennett, M.B.		
Report Card Remarks	Mostly discarded when caught in fisheries, with generally stable catch rates		

### Summary

The Australian Sawtail Shark is a small apparently common, catshark that is widespread across southern Australia. The species is retained as byproduct and is also a regular discarded bycatch in various demersal trawl fisheries. A minor decline in catch rates has been documented in heavily fished areas of southeast Australia, while elsewhere catch rates have been stable. An oviparous species, it is likely to be productive, however catch rates should be monitored in heavily fished areas. It has a wide bathymetric range across the outer continental shelf and upper slope, and likely finds refuge in unfished areas. Therefore, the species is assessed as Least Concern (IUCN) and Sustainable (SAFS).



### Distribution

The Australian Sawtail Shark occurs in temperate to subtropical waters, ranging from Noosa, southeast Queensland to Carnarvon, Western Australia, and including the waters of New South Wales, Victoria, Tasmania and South Australia (Compagno and Niem 1998, Last and Stevens 2009).

### Stock structure and status

The Australian Sawtail Shark is reportedly common (Walker and Gason 2007, Last and Stevens 2009). Catch rates recorded in two regions covered by this fishery were stable in one (southern Australia; 2000-2006), and showed a minor but significant decline in the other (southeast Australia; 1988-2006). In bycatch surveys of the eastern king prawn sector (deepwater component) of the Queensland East Coast Trawl Fishery, this shark was the third most commonly caught elasmobranch, although numbers were low and survivorship from trawling was high (Courtney et al. 2014). Post-release mortality is unknown.

## Fisheries

The species is not targeted by commercial fisheries and is only of minor importance to fisheries through retention as byproduct in the Western Australian Deepwater Trawl Fishery, although the quantity is unknown (Rose and SAG 2001). It is a frequent component of discarded bycatch in other demersal trawl fisheries across its range. It is discarded in the Southern and Eastern Scalefish and Shark Fishery where it is regularly caught by trawlers (Walker and Gason 2007).

## Habitat and biology

The Australian Sawtail Shark is demersal on the outer continental shelf and upper slope at depths of 130 to 640 m. Maximum size is 61 cm total length (TL), with both sexes maturing at approximately 40 cm TL (Last and Stevens 2009, Kyne et al. 2011). It appears to sometimes aggregate by sex (Last and Stevens 2009).

Longevity and maximum size	Longevity: unknown Max size: 61 cm TL
Age and/or size at maturity (50%)	Both sexes: ~40 cm TL

**Link to IUCN Page:** <http://www.iucnredlist.org/details/41811/0>

**Link to page at Shark References:** <http://www.shark-references.com/species/view/Figaro-boardmani>

## References

- Compagno, L.J.V. and Niem, V.H. 1998. Scyliorhinidae. In: K.E., Carpenter and V.H., Niem (eds) FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 2. Cephalopods, crustaceans, holothurians and sharks. FAO, Rome
- Courtney, A.J., Campbell, M.J., Tonks, M.L., Roy, D.P., Gaddes, S.W., Haddy, J.A., Kyne, P.M., Mayer, D.G. and Chilcott, K.E. 2014. Effects of bycatch reduction devices in Queensland's (Australia) deepwater eastern king prawn (*Melicertus plebejus*) trawl fishery. *Fisheries Research* 157(2014): 113-123.
- Kyne, P.M., Courtney, A.J. and Bennett, M.B. 2011. Observations on the reproductive biology of three catsharks (Carcharhiniformes: Scyliorhinidae: *Asymbolus* and *Figaro*) from the continental shelf of southern Queensland, Australia. *Journal of the Marine Biological Association of the United Kingdom* 91: 1157-1164.
- Last, P.R. and Stevens, J.D. 2009. Sharks and Rays of Australia. Second Edition. CSIRO Publishing, Collingwood.
- Rose, C. and Shark Advisory Group (SAG). 2001. Australian Shark Assessment Report. Commonwealth Department of Agriculture, Fisheries and Forestry – Australia, Canberra.
- Walker, T.I. and Gason, A.S. 2007. Shark and other chondrichthyan byproduct and bycatch estimation in the Southern and Eastern Scalefish and Shark Fishery. Final report to Fisheries and Research Development Corporation Project No. 2001/007. July 2007. vi + 182 pp. Primary Industries Research Victoria, Queenscliff, Victoria, Australia.