

## Bluntnose Sixgill Shark, *Hexanchus griseus*

Report Card assessment	Sustainable		
IUCN Red List Australian Assessment	Refer to Global Assessment	IUCN Red List Global Assessment	Near Threatened
Assessors	Barnett, A. & Braccini, J.M.		
Report Card Remarks	Rare in Australia with catches monitored to ensure they remain within acceptable levels		

### Summary

The Bluntnose Sixgill Shark is a wide ranging but patchily distributed species in boreal, temperate and tropical waters. It is taken as bycatch in a wide range of fishing gear and is considered a valuable food and sports fish. In Australia, it is caught in southeastern fisheries and has been classed as at high ecological risk and its capture in fisheries is monitored to ensure catches remain low. The species appears vulnerable to overfishing and unable to sustain intensive, targeted fisheries for long periods. Population and fisheries data, however, are lacking from many regions. In Australian waters, it is considered rare with catches monitored to ensure they remain within acceptable levels; there is no indication that the population has declined to levels where it would be threatened. Therefore, the species is assessed as Near Threatened (IUCN) and in Australia, Sustainable (SAFS).



### Distribution

The species is widely but patchily distributed across the Pacific, Atlantic and Indian Oceans. In Australia, it is recorded from seamounts off Queensland and from New South Wales, Victoria, Tasmania and Western Australia (Last and Stevens 2009).

### Stock structure and status

There is currently very little information on population size, structure, or trend for the species

### Fisheries

Due to its broad depth range and relative sluggishness, the species is often caught as bycatch of longline, handline, gillnet, traps, trammel net and pelagic and bottom trawls (Clarke et al. 2005, Fowler et al. 1997, Arocha et al. 2002, Walker et al. 2008). It has been commercially utilised, including in Australia, for meat and liver oil. In southeast Australia, it is considered rare based on reported catches, and to have moderate to low catch susceptibility to trawl, gillnet and hook fishing gears. It is classed as high ecological risk in terms of abundance and moderate risk in terms of catch susceptibility (Walker et al. 2008) and the levels of catch in fisheries are monitored to ensure they remain within acceptable levels.

## Habitat and biology

The Bluntnose Sixgill Shark occurs along the outer shelves and upper slopes from the surface to approximately 2,500 m (Last and Stevens 2009). It occasionally comes close inshore in areas adjacent to deepwater trenches or shelves (Dunbrack 2008, Andrews et al. 2009). In Tasmania, a single individual was caught approximately 30 km up the Derwent River, over 100 km from the deeper waters of the continental shelf (Barnett et al. 2010). Maximum size is at least 482 cm total length (TL) with males mature at 315 cm TL and females mature at >400 cm TL (Last and Stevens 2009).

Longevity and maximum size	Longevity unknown Max size: at least 482 cm TL
Age and/or size at maturity (50%)	Males: 315 cm TL Females: >400 cm TL

**Link to IUCN Page:** <http://www.iucnredlist.org/details/10030/0>

**Link to page at Shark References:** <http://www.shark-references.com/species/view/Hexanchus-griseus>

### References

- Arocha F., Arocha O. & Marcano L. 2002. Observed shark bycatch from the Venezuelan tuna and swordfish fishery from 1994 through 2000. ICCAT Collective Volume of Scientific Papers. Andrews KS, Williams GD, Farrer D, Tolimieri N, Harvey C J, Bargmann G & Levin PS. 2009. Diel activity patterns of sixgill sharks, *Hexanchus griseus*: the ups and downs of an apex predator. *Animal Behaviour* 78(525–536.).
- Andrews K.S., Williams G.D., Farrer D., Tolimieri N., Harvey C.J., Bargmann G. & Levin P.S. 2009. Diel activity patterns of sixgill sharks, *Hexanchus griseus*: the ups and downs of an apex predator. *Animal Behaviour* 78(525–536.).
- Barnett A., Stevens J.D. & Yick J.L. 2010. The occurrence of the bluntnose sixgill shark *Hexanchus griseus* (Hexanchiformes: Hexanchidae) in a river in south-eastern Tasmania. *Marine Biodiversity Records* 3: e24.
- Clarke, M., Borges, L., Officer, R. and Stokes, D. 2005. Comparisons of Trawl and Longline Catches of Deepwater Elasmobranchs West and North of Ireland. *Journal of Northwest Atlantic Fisheries Science* 35: 429-442.
- COSEWIC. 2007. Assessment and Status Report on the Bluntnose Sixgill Shark *Hexanchus griseus* in Canada. In: Committee on the Status of Endangered Wildlife in Canada (ed.) Ottawa.
- Dunbrack R. 2008. Abundance trends for *Hexanchus griseus* (bluntnose sixgill shark) and *Hydrolagus colliciei* (spotted ratfish) counted at an automated underwater observation station in the Strait of Georgia, British Columbia. *Canadian Field-Naturalist* 122: 124–128.
- Fowler S.L., Reed T.M. & Dipper F.A. 1997. Elasmobranch biodiversity, conservation and management. In: Fowler S.L., Reed T.M. & Dipper F.A. (ed.), International Seminar and Workshop, pp. 9–13. Sabah, Malaysia.
- Larson S., Christiansen J., Griffing D., Ashe J., Lowry D. & Andrews K. 2010. Relatedness and polyandry of sixgill sharks, *Hexanchus griseus*, in an urban estuary. *Conservation Genetics* 12: 679–690.
- Last, P. R. and J. D. Stevens 2009. *Sharks and Rays of Australia*, Second Edition. Collingwood, Australia, CSIRO Publishing.
- McFarlane, G. A., King, J. R. & Saunders, M. W. 2002. Preliminary study on the use of neural arches in the age determination of bluntnose sixgill sharks (*Hexanchus griseus*). *Fishery Bulletin* 100: 861–864.
- Walker T.I., Stevens J.D., Braccini J.M., Daley R.K., Huveneers C., Irvine S.B., Bell J.D., Tovar-Avila J., Trinnie F.I., Phillips D.T., Treloar M.A., Awruch C.A., Gason A.S., Salini J. & Hamlett W.C. 2008. Rapid Assessment of Sustainability for Ecological Risk of Shark and Other Chondrichthyan Bycatch Species Taken in the Southern and Eastern Scalefish and Shark Fishery. Primary Industries Research Victoria, Queenscliff.
- Williams G.D., Andrews K.S., Farrer D.A. & Levin P.S. 2010. Catch rates and biological characteristics of bluntnose sixgill sharks in Puget Sound. *Transactions of the American Fisheries Society* 139: 108–116.