

## Broadnose Sevengill Shark, *Notorynchus cepedianus*

Report Card assessment	Sustainable		
IUCN Red List Australian Assessment	Least Concern	IUCN Red List Global Assessment	Data Deficient
Assessors	Barnett, A. J. & Braccini, M.		
Report Card Remarks	Limited catch data suggests Australian stock increasing after previous years of reduction		

### Summary

The Broadnose Sevengill Shark is a wide ranging and moderately common, temperate water species. It is exposed to intensive inshore fisheries over most of its range. In Australia, the species is a bycatch in the southern trawl, gillnet and longline fisheries. It is also taken by recreational anglers. Limited fisheries catch data suggests the Australia stock may be increasing after previous years of reduction due to the introduction of gear restrictions. Elsewhere there is a lack of population and catch data.



Source: Ross Robertson / Shorefishes of the tropical eastern Pacific online information system. License: CC BY Attribution

Therefore, globally it is assessed as Data Deficient (IUCN), and in Australia it is assessed as Least Concern (IUCN) and Sustainable (SAFS).

### Distribution

The Broadnose Sevengill Shark occurs in temperate coastal waters of the south Atlantic, Pacific and Indian Oceans. In Australia, it is found in southern waters from Sydney (New South Wales) to Esperance (Western Australia) (Last and Stevens 2009).

### Stock structure and status

There is currently limited information on population size or trend for the species. The disjunct worldwide distribution of this species suggests subpopulations. Genetic data reveal significant differentiation between samples from the west Pacific, eastern North Pacific and south Atlantic, indicating the presence of at least three genetically distinct populations. In Australia, movement information suggests a single population (Barnett et al. 2011, Stehfest et al. 2014). The limited Australian catch data suggests the stock may be increasing after previous years of reduction, due to the introduction of gear restrictions (Walker and Gason, 2007, Braccini et al. 2009).

## Fisheries

The Broadnose Sevengill Shark is taken as bycatch of intensive inshore fisheries and sometimes targeted in commercial and recreational fisheries over most of its range. The flesh is good quality and the hide and liver oil are also used. In southeast Australia, the species is commonly taken as bycatch in the trawl, gillnet and longline catches of the Southern and Eastern Scalefish and Shark Fishery (Walker et al. 2005, Walker and Gason 2009). The species has a moderate to low catch susceptibility to trawl and hook fishing gears, but high vulnerability to gillnets, categorizing the species as at high ecological risk in terms of abundance and high risk in terms of catch susceptibility (Walker et al. 2007). The species was also considered at high risk in the gillnet sector in a separate sustainability assessment (Zhou et al. 2007) and catches are monitored to ensure that catches remain low. Globally, there is some catch data, however the quality of the data is uncertain.

## Habitat and biology

The Broadnose Sevengill Shark occur from close inshore, in bays and estuaries, out to at least 360 m on the edge of continental shelves (Stehfest et al. 2014). Maximum size is 300 cm total length (TL) and maximum age is 32 years. Males mature at 4-5 years and 150 cm TL and females at 11-21 years and 220 cm TL (Van Dykhuizen and Mollet 1992, Braccini et al. 2010).

Longevity and maximum size	Longevity: 32 years Max size: 300 cm TL
Age and/or size at maturity (50%)	Males: 4-5 years, 150 cm TL Females: 11-21 years, 220 cm TL

**Link to IUCN Page:** <http://www.iucnredlist.org/details/39324/0>

**Link to page at Shark References:** <http://www.shark-references.com/species/view/Notorynchus-cepedianus>

## References

- Barnett, A., Abrantes K.G., Stevens J.D. and Semmens J.M. 2011. Site fidelity and sexspecific migration in a mobile apex predator: implications for conservation and ecosystem dynamics. *Animal Behaviour* 81: 1039–1048.
- Braccini J.M., Walker, T.I., and Gason, A.S. 2009. GHATF shark survey of population abundance and population size composition for target, byproduct and bycatch species. Report to Australian Fisheries Management Authority. June 2009. iv + 123 pp. Fisheries Research Branch, Department of Primary Industries, Queenscliff, Victoria, Australia.
- Braccini J.M., Troynikov V.S., Walker T.I., Mollet H.F., Ebert D.A., Barnett A. and Kirby N. 2010. Embracing growth heterogeneity of wild and captive broadnose sevengill sharks, *Notorynchus cepedianus*, at a global scale. *Aquatic Biology* 9: 131–138.
- Last, P.R. and Stevens, J.D. 2009. Sharks and Rays of Australia, 2nd edition. CSIRO, Melbourne, Australia.
- Stehfest K.M., Patterson T.A., Barnett A. and Semmens J.M. 2014. Intraspecific differences in movement, dive behaviour and vertical habitat preferences of a key marine apex predator. *Marine Ecology Progress Series* 495: 249-262.
- Van Dykhuizen, G. and Mollet, H.F. 1992. Growth, age estimation, and feeding of captive sevengill sharks, *Notorynchus cepedianus*, at the Monterey Bay Aquarium. In: J.G. Pepperell (ed.), Sharks: Biology and fisheries. *Australian Journal of Marine and Freshwater Research*, pp. 297–318.
- Walker T., Dowdney J., Williams A., Fuller M., Webb H., Bulman C., Sporcic M. and Wayte S. 2007. Ecological Risk Assessment for the Effects of Fishing: Report for the Shark Gillnet Component of the Gillnet Hook and Trap Sector of the Southern and Eastern Scalefish and Shark Fishery. Australian Fisheries Management, Canberra.
- Walker T.I. and Gason A.S. 2009. SESSF monitoring data management, reporting and documentation 2006/07. Australian Fisheries Management Authority, Canberra.
- Walker T.I., Hudson R.J. and Gason A.S. 2005. Catch evaluation of target, by-product and by-catch species taken by gillnets and longlines in the shark fishery of south-eastern Australia. *Journal of Northwest Atlantic Fishery Science* 35: 505–530.
- Walker, T.I. and Gason, A.S. 2007. Shark and other chondrichthyan byproduct and bycatch estimation in the Southern and Eastern Scalefish and Shark Fishery. Final report to Fisheries Research and Development Corporation Project No. 2001/007. (July 2007) pp. 182 + vi. Primary Industries Research Victoria: Queenscliff, Victoria, Australia, Victoria.
- Zhou S., Smith A. and Fuller M. 2007. Rapid Quantitative Risk Assessment for Fish Species in Selected Commonwealth Fisheries. Australian Fisheries Management Authority, Canberra.