

Great Hammerhead, *Sphyrna mokarran*

Report Card assessment	Depleted		
IUCN Red List Australian Assessment	Vulnerable	IUCN Red List Global Assessment	Endangered
Assessors	Heupel, M.R.		
Report Card Remarks	In Australia, considered potentially at high risk; globally in decline		

Summary

The Great Hammerhead is a large, globally distributed tropical shark. The species is highly valued for its fins and is taken in a large range of coastal fisheries. It has very high bycatch mortality and only reproduces once every two years, making it vulnerable to over-exploitation and population depletion. In Australia, capture in commercial and recreational fisheries, genetic links and connectivity with Indonesian populations, intense fishing pressure in Indonesian waters and potentially IUU fishing all contribute to declines in the Australian stock. Risk assessments of northern Australian elasmobranchs indicate that Great Hammerhead may be 'high-risk'. However, species specific data on population status is limited. Further study is urgently required to more fully determine the status of Great Hammerhead in Australia. Given the decline in global populations, its vulnerability to depletion, low survival at capture and high value for the fin trade, the species is assessed as globally Endangered (IUCN). In Australia, the species is assessed as Vulnerable (IUCN) and Overfished (SAFS).



Risk assessments of northern Australian elasmobranchs indicate that Great Hammerhead may be 'high-risk'. However, species specific data on population status is limited. Further study is urgently required to more fully determine the status of Great Hammerhead in Australia. Given the decline in global populations, its vulnerability to depletion, low survival at capture and high value for the fin trade, the species is assessed as globally Endangered (IUCN). In Australia, the species is assessed as Vulnerable (IUCN) and Overfished (SAFS).

Distribution

The Great Hammerhead ranges widely throughout the tropical and warm temperate waters of the world (Last and Stevens 2009). In Australia, it occurs from Sydney (New South Wales) across the tropics to Mandurah (Western Australia) (Last and Stevens 2009).

Stock structure and status

Little is known about the population size and trend of the Great Hammerhead. Generally regarded as solitary, the species is therefore unlikely to be abundant wherever it occurs. The closely related and similarly distributed Scalloped Hammerhead shows genetic structuring between ocean basins with little structuring evident within basins (Duncan et al. 2006). Genetic connectivity between Australia and Indonesia is possible for the Great Hammerhead, given its potential for pelagic movement but needs to be confirmed. Species specific data on population status in Australian waters is limited, partly because of confusion with Scalloped Hammerheads. Declines within Australian populations have been

reported for Scalloped Hammerhead populations on both the east and west coast and declines are also presumed for Great Hammerhead although direct data are lacking. As such, the population is inferred to be Vulnerable (IUCN) and Overfished (SAFS). Further study is urgently required to more fully determine the status of Great Hammerhead in Australia.

Fisheries

The primary threat to the Great Hammerhead is fishing. The dorsal fin is particularly large and highly valued and the species is taken by target and bycatch fisheries (Dudley and Simpfendorfer 2006, Zeeberg et al. 2006). It is regularly caught in the tropics with longlines, fixed bottom nets, hook-and-line, and possibly with pelagic and bottom trawls. In Australia, there was a directed shark fishery operated by Taiwan around the northern coast of Australia that regularly caught Great Hammerheads up until 1986 (Stevens and Lyle 1989). Illegal, unregulated and unreported (IUU) fishing has been significant in northern Australia. Increased foreign fishing was observed between 2001 and 2005 with Indonesian vessels joining Chinese and Taiwanese fleets exploiting this region (Field et al. 2009). Small scale vessels were estimated to have harvested between 300 and 1,100 tonnes of shark in 2006 (Marshall 2011). Industrial scale Taiwanese boats were estimated to have harvested up to 100 times that amount. There has been a decline since 2006 in IUU fishing in northern Australian waters (Field et al. 2009, Haward and Bergin 2016), however it is still likely to occur. The Great Hammerhead continues to be taken in multiple fisheries including gillnet, trawl, line fisheries although little species-specific data are available. Other threats include sport fishing (Pepperell 1992) and capture in shark control programs in New South Wales and Queensland.

Fishing in neighbouring countries may also have implications for Great Hammerhead in Australia, given it is likely to be a shared stock. Indonesia is one of the world's largest fishers of sharks with annual catch rates estimated at 110,000 tonnes, although there is growing evidence that the catch rates are declining (White et al. 2006, Blaber et al. 2009, Lack and Sant 2009). As localised depletions become apparent, fishers are moving to exploit new areas (White and Kyne 2010). Intense fishing pressure in southeast Asia has led to significant depletions in coastal fisheries resources that includes sharks (Stobutzki et al. 2006).

Habitat and biology

The Great Hammerhead is a coastal-pelagic and semi-oceanic species that occurs close inshore and well offshore at depths ranging from near surface to over 80 m (Compagno 1984). Maximum size is 550 to 610 cm total length (TL), though 400 cm TL is more common for a mature adult (Compagno 1984, Last and Stevens 2009). Maximum age is estimated at 39 years (Harry et al. 2011). Males mature at approximately 228-269 cm TL and females mature at 228 to 300 cm TL (Compagno 1984) with both sexes mature at 8 years (Harry et al. 2011). Litter size range from 6 to 42 pups (Last and Stevens 2009) and females breed once every two years (Stevens and Lyle 1989).

Longevity and maximum size	Longevity: estimated 39 years Max size: 550-610 cm TL
Age and/or size at maturity (50%)	Males: 8 years, 228-269 cm TL Females: 8 years, 228-300 cm TL

Link to IUCN Page: <http://www.iucnredlist.org/details/39386/0>

Link to page at Shark References: <http://www.shark-references.com/species/view/Sphyrna-mokarran>

References

- Blaber, S., Dichmont, C.M., White, W.T., Buckworth, R.C., Sadiyah, L., Iskandar, B., Nurhakim, S., Pillans, R.D., Andamari, R., Dharmadi and Fahmi. 2009. Elasmobranchs in southern Indonesian fisheries: the fisheries, the status of the stocks and management options. *Reviews in Fish Biology and Fisheries* 19: 367–391.
- Compagno, L.J.V. 1984. FAO Species Catalogue. Sharks of the World: an annotated and illustrated catalogue of the shark species known to date. Part 2 - Carcharhiniformes. FAO Fisheries Synopsis No. 125, Vol. 4(2). FAO, Rome.
- Dudley, S.F.J. and Simpfendorfer, C.A. 2006. Population status of 14 shark species caught in the protective gillnets off KwaZulu-Natal beaches, South Africa, 1978-2003. *Marine and Freshwater Research* 57: 225-240.
- Duncan K., Martin A., Bowen B. W. and De Couet H. G. 2006. Global phylogeography of the scalloped hammerhead shark (*Sphyrna lewini*). *Molecular Ecology* 15: 2239-2251.
- Field, I.C., Meekan, M.G., Buckworth, R.C. and Bradshaw, J.A. 2009. Protein mining the world's oceans. Australasia as an example of illegal expansion-and-displacement fishing. *Fish and Fisheries* 10: 323-328.
- Harry A. V., Macbeth W. G., Gutteridge A. N. and Simpfendorfer C. A. 2011. The life histories of endangered hammerhead sharks (Carcharhiniformes, Sphyrnidae) from the east coast of Australia. *Journal of Fish Biology* 78: 2026-2051.
- Haward, M. and Bergin, A. 2016. Net worth: Australia's regional fisheries engagement. Australian Strategic Policy Unit, Canberra.
- Lack, M. and Sant, G. 2009. Trends in Global Shark Catch and Recent Developments in Management. TRAFFIC International, Cambridge, UK.
- Last, P.R. and Stevens, J.D. 2009. Sharks and Rays of Australia. Second Edition. CSIRO Publishing, Collingwood.
- Marshall, L.J. 2011. The Fin Blue Line: Quantifying fishing mortality using shark fin morphology. PhD Thesis. University of Tasmania.
- Ovenden, J.R., Kashiwagi, T., Broderick, D., Giles, J., and Salini, J. 2009. The extent of population genetic subdivision differs among four co-distributed shark species in the Indo-Australian archipelago. *BMC Evolutionary Biology* 9(40).
- Ovenden J. R., Morgan J. A. T., Street R., Tobin A., Simpfendorfer C., Macbeth W. and Welch D. 2011. Negligible evidence for regional genetic population structure for two shark species *Rhizoprionodon acutus* (Ruppell, 1837) and *Sphyrna lewini* (Griffith & Smith, 1834) with contrasting biology. *Marine Biology* 158: 1497-1509.
- Pepperell, J.G. 1992. Trends in the distribution, species composition and size of sharks caught by gamefish anglers off south-eastern Australia, 1961-90. *Australian Journal of Marine and Freshwater Research* 43: 213-25
- Stevens, J.D. and Lyle, J.M. 1989. Biology of three hammerhead sharks (*Eusphyra blochii*, *Sphyrna mokarran* and *S. lewini*) from Northern Australia. *Australian Journal of Marine and Freshwater Research* 40:129–146.
- Stobutzki I. C., Silvestre G. T. and Garces L. R. 2006. Key issues in coastal fisheries in South and Southeast Asia, outcomes of a regional initiative. *Fisheries Research* 78: 109-118.
- White, W.T., Last, P.R., Stevens, J.D., Yearsley, G.K., Fahmi and Dharmadi. 2006. Economically Important Sharks and Rays of Indonesia. Australian Centre for International Agricultural Research, Canberra, Australia.
- White, W.T. and Kyne P.M. 2010. The status of chondrichthyan conservation in the Indo-Australasian region. *Journal of Fish Biology* 76: 2090-2117.
- Zeeberg, J., Coorten, A. and Graaf, E. 2006. Bycatch and release of pelagic megafauna in industrial trawler fisheries off Northwest Africa. *Fisheries Research* 78: 186-195.