

Southern Bluefin Tuna (2018)

Thunnus maccoyii



Heather Patterson: Australian Bureau of Agricultural and Resource Economics and Sciences, **Simon Nicol:** Australian Bureau of Agricultural and Resource Economics and Sciences

STOCK STATUS OVERVIEW

Jurisdiction	Stock	Fisheries	Stock status	Indicators
Commonwealth	Global	CCSBT, SBTF	Recovering	Spawning stock biomass, projections of rebuilding

CCSBT Commission for the Conservation of Southern Bluefin Tuna (CTH), SBTF Southern Bluefin Tuna Fishery (CTH)

STOCK STRUCTURE

Southern Bluefin Tuna constitutes a single, highly migratory biological stock that spawns in the north-east Indian Ocean and migrates throughout the temperate southern oceans, supporting a number of international fisheries [Evans et al. 2012, Patterson et al. 2018, Proctor et al. 1995].

Here, assessment of stock status is presented at the biological stock level—Global.

STOCK STATUS

Global The biological stock of Southern Bluefin Tuna is fished by Australian fishers endorsed to fish in the Southern Bluefin Tuna Fishery (Commonwealth) —and members of the Commission for the Conservation of Southern Bluefin Tuna. Fishers in the Eastern Tuna and Billfish Fishery (Commonwealth) and the Western Tuna and Billfish Fishery (Commonwealth) who have quota for Southern Bluefin Tuna can retain catches of the species. The species is also caught by recreational fishers in the waters off southern Australia, and there are other sources of unaccounted mortalities, including catches by fleets that are not members of the Commission for the Conservation of Southern Bluefin Tuna (CCSBT) [CCSBT 2015, Patterson et al. 2018].

In 2011, the CCSBT adopted a management procedure (analogous to a harvest strategy, evaluated fully using management strategy evaluation) to guide the recovery of the biological stock to 20 per cent of unfished biomass by 2035 with 70 per cent probability [Hillary et al. 2016]. Performance of the management procedure is monitored using the biomass of fish that are 10 years and older. The most recent assessment (2017) undertaken by the CCSBT takes into account reported catch from all international jurisdictions [CCSBT 2017]. It also examines the sensitivity of the results to alternate scenarios for unaccounted

fishing mortalities. The actual (current and historic) level of unaccounted fishing mortality from all sources is uncertain, but there are indications it may be substantial [Patterson et al. 2018].

The most recent assessment estimated that the biomass of fish 10 years and older in the Southern Bluefin Tuna biological stock is still low, at 11 per cent of unfished levels (9 to 13 per cent across reference set of models) [CCSBT 2017, Hillary et al. 2017]. This is a progressive increase since the 2014 assessment which was 7 per cent [CCSBT 2017]. The most recent estimate of spawning stock biomass is 13 per cent (11 to 17 per cent across reference set of models) of unfished levels [CCSBT 2017]. The assessment also indicated that the stock may be more productive than for previous assessments and recent recruitment levels are estimated to be above long-term average levels. Based on the estimated stock size (13 per cent), the stock is considered to be recruitment impaired.

The estimate of fishing mortality was 50 per cent (38 to 66 per cent for reference set of models) of the level associated with maximum sustainable yield. Although there may be substantial unaccounted mortality of Southern Bluefin Tuna [CCSBT 2017], projections of the performance of the management procedure under scenarios of different levels of unaccounted mortalities showed that these mortalities are unlikely to reduce the probability of rebuilding to the specified interim management target to below the required 70 per cent within the specified time frame [CCSBT 2017]. In fact, the projections indicate that the stock may recover to 20 per cent of unfished levels before 2035.

On the basis of the evidence provided above, the global biological stock is classified as a **recovering stock**.

BIOLOGY

Southern Bluefin Tuna biology [David et al. 2001, Farley et al. 2014, 2015, Laslett et al. 2002]

Species	Longevity / Maximum Size	Maturity (50 per cent)
Southern Bluefin Tuna	40+ years; ~1 900 mm FL	~10–12 years; 1 580–1 630 mm FL

DISTRIBUTION



Distribution of reported commercial catch of Southern Bluefin Tuna

TABLES

Commercial Catch Methods	Commonwealth
Gillnet	✓
Hand Line, Hand Reel or Powered Reels	✓
Handline	✓
Pelagic Longline	✓
Pole and Line	✓
Purse Seine	✓
Trolling	✓
Unspecified	✓
Various	✓

Fishing methods	Commonwealth
Commercial	
Gillnet	✓
Handline	✓
Pelagic Longline	✓
Pole and Line	✓
Purse Seine	✓
Trolling	✓
Various	✓
Recreational	

Hook and Line	✓
Spearfishing	✓
Management Methods	
	Commonwealth
Commercial	
Area restrictions	✓
Catch limits	✓
Individual transferable quota	✓
Recreational	
Bag limits	✓
Boat limits	✓
Active Vessels	
	Commonwealth
	15 Vessels in ETBF, 1 Vessels in WTBF, 11 Vessels in SBTF,

ETBF Eastern Tuna and Billfish Fishery(CTH)

WTBF Western Tuna Billfish Fishery(CTH)

SBTF Southern Bluefin Tuna Fishery(CTH)

Catch	
	Commonwealth
Commercial	8613.27t in CCSBT, 5334t in SBTF,
Indigenous	Unknown
Recreational	Unknown

CCSBT Commission for the Conservation of Southern Bluefin Tuna (CTH), SBTF Southern Bluefin Tuna Fishery (CTH),


Commonwealth – Recreational The Australian Government does not manage recreational fishing in Commonwealth waters. Recreational fishing in Commonwealth waters is managed by the state or territory immediately adjacent to those waters, under its management regulations. Recreational catches reported here are from state surveys during specific time periods, as noted.

Commonwealth Recreational and Indigenous – Recreational and Indigenous fishing sectors reported here are New South Wales, Queensland, South Australia, Tasmania, Victoria and Western Australia. Measures listed here exist in at least one of these jurisdictions.

Commonwealth – Indigenous The Australian Government does not manage non-commercial Indigenous fishing in Commonwealth waters, with the exception of the Torres Strait. In general, non-commercial Indigenous fishing in Commonwealth waters is managed by the state or territory immediately adjacent to those waters.

Commonwealth – Commercial (catch) Catches reported for the Commission for the Conservation of Southern Bluefin Tuna (CCSBT) are for 2016, the most recent year available.

CATCH CHART



Commercial catch of Southern Bluefin Tuna - note confidential catch not shown

EFFECTS OF FISHING ON THE MARINE ENVIRONMENT

ENVIRONMENTAL EFFECTS on Southern Bluefin Tuna

References	
710	Evans, k, Patterson, TA, Reid, H and Harley, SJ 2012, Reproductive schedules in southern bluefin tuna: Are current assumptions appropriate? PLoS One, 7(4):e34550.
711	Patterson, TA, Eveson, JP, Hartog, JR, Evans, K, Cooper, S, Lansdell, M, Hobday, AJ and Davies, CJ 2018, Migration dynamics of juvenile southern bluefin tuna, Scientific Reports, 8: 14553.
712	Farley, JH, Eveson, JP, Davis, TLO, Andamari, R, Proctor, CH, Nugraha, B and Davies, CR 2014, Demographic structure, sex ration and growth rates of southern bluefin tuna (<i>Thunnus maccoyii</i>) on the spawning ground, PLoS One 9(5): e96392.
713	Farley, JH, David, TLO, Bravington, MV, Andamari, R and Davies, CR 2015, Spawning dynamics and size related trends in reproductive parameters of southern bluefin tuna, <i>Thunnus maccoyii</i> , PLoS One 10(5): e0125744.
714	Hillary, RM, Preece, AL, Davies, CR, Kurota, H, Sakai, O, Itoh, T, Parma, AM, Butterworth, DS, Ianelli, J and Branch, TA 2016, A scientific alternative tomoratoria for rebuilding depleted international tuna stocks, <i>Fish and Fisheries</i> , 17(2): 469–482.
715	Hillary, RM, Preece, AL and Davies CR 2017, Reconditioning of the CCSBT operating model in 2017, paper CCSBT-ESC/1708/14 for the twenty second meeting of the CCSBT Scientific Committee, 28 August–2 September 2017, Yogyakarta, Indonesia.
716	Proctor, CH, Thresher, RE, Gunn, JS, Mills, DJ, Harrowfield, IR and Sie, SH 1995, Stock structure of the Southern Bluefin Tuna, <i>Thunnus maccoyii</i> : an investigation based on probe micro analysis of otolith composition, <i>Marine Biology</i> , 122(4): 511–526.
717	Commission for the Conservation of Southern Bluefin Tuna 2015, Report of the twentieth meeting of the Scientific Committee, Incheon, South Korea, 1–5 September 2015.
718	Patterson, H, Nicol, S and Curtotti, R 2018, Southern Bluefin Tuna Fishery, in H Patterson, J Larcombe, S Nicol and R Curtotti (eds), <i>Fishery status reports 2018</i> , Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra, 393–403.
719	Commission for the Conservation of Southern Bluefin Tuna 2017, Report of the twenty second meeting of the Scientific Committee, Yogyakarta, Indonesia, 28 August–2 September 2017.
720	Davis, T, Farley, J and Gunn, J 2001, Size and age at 50% maturity in SBT: an integrated view from published information and new data from the spawning ground, CCSBT-

	SC/0108/16, Tokyo, Japan, 28–31 August 2001.
721	Laslett, GM, Eveson, JP and Polacheck, T 2002, A flexible maximum likelihood approach for fitting growth curves to tag-recapture data, Canadian Journal of Fisheries and Aquatic Sciences, 59: 976–986.