

## Southern Dogfish, *Centrophorus uyato*

<b>Report Card assessment</b>	<b>Depleted</b>		
IUCN Red List Australian Assessment	Endangered	IUCN Red List Global Assessment	Endangered
Assessors	Finucci, B., Bineesh, K.K., Cotton, C.F., Dharmadi, Kulka, D.W., Neat, F.C., Pacoureau, N., Rigby, C.L., Tanaka, S. & Walker, T.I.		
Australian Assessors	Kyne, P.M., Heupel, M.R., White, W.T. & Simpfendorfer, C.A. (Shark Action Plan)		
Report Card Remarks	In Australia, severely depleted with management measures implemented but recovery expected to be slow. Listed on EPBC Act (Conservation Dependent).		

### Summary

The Southern Dogfish is a deepwater species inhabiting continental slopes of Australia, Africa, Europe and Asia. It was formally unedified in Australia as *Centrophorus zeehani*, but taxonomic revision resulted in it being aligned with the more widespread *C. uyato*. It is taken as bycatch by deepwater fisheries and depletion estimates for this dogfish off eastern Australia indicated a 79% reduction in virgin population size. This part of its range is still fished and there is no evidence that numbers have recovered on these fishing grounds. It has very low productivity that make the species extremely sensitive to rapid population depletion by commercial fishing and also means very slow recovery after such depletion. A number of conservation measures have been implemented to promote recovery following its listing as Conservation Dependent (EPBC Act). These include a total ban on retaining any specimens for sale, gulper shark (*Centrophorus* spp.) protection areas off Sydney and eastern Bass Strait that are closed to all methods of fishing, and a ban on trawling below 700 m. Therefore, the species is assessed as Endangered (IUCN) in Australia (Kyne et al. 2021) and Depleted (SAFS) because although management is in place, recovery has not yet been shown and is expected to take decades.



### Distribution

Southern Dogfish occurs off southern Australia, patchily through Asia, and in the eastern Atlantic (including Africa and Europe, and the western North Atlantic and Caribbean (White et al 2022). In Australia, it is found from southern Queensland to Shark Bay.

### Stock structure and status

Southern Dogfish has been depleted off New South Wales between Sydney and the Eden-Gabo Island area where over 20 years from 1976–1977 to 1996–1997 there was a >99% decline in relative

abundance of all gulper sharks (*Centrophorus* spp.) (Andrew et al. 1997, Graham et al. 2001). It has also historically been fished in South Australia and Western Australia, but information on catches and stock status are unavailable.

## Fisheries

The core depth of the species (350–800 m) coincides with the most heavily fished depths by trawlers and longliners operating on the upper slope around southeast Australia. In the period 1975–2000, the population south of Newcastle (NSW) was severely affected with its relative abundance reduced to <5% of historical levels (Graham et al. 2001, Daley et al. 2002, Wilson et al. 2009). Commercial fishing in the Commonwealth Trawl and Scalefish Hook Sectors of the Southern and Eastern Scalefish and Shark Fishery (SESSF) continued to affect the remaining stock, despite a ban on trawling below 700 m implemented in 2007 (AFMA 2006). In 2010, the stock status for the three species of upper slope gulper sharks (Harrison’s Dogfish, Endeavour Dogfish *C. moluccensis* and Southern Dogfish *C. uyato/C. zeehaani*) on southeast Australian grounds was assessed as ‘overfished’ and ‘subject to overfishing’ (Stobutzki et al. 2011). Further south off eastern Bass Strait and Tasmania, trawling and targeted gillnet fishing in the 1980s and 1990s also severely depleted numbers (Daley et al. 2002). As a result of population declines a plan of management was implemented (AFMA 2012) and the species was listed as Conservation Dependent under the *Environment Protection Biodiversity Conservation Act 1999* (EPBC Act) in 2013. This plan includes closures to allow the recovery of the population, no retention and other spatial and temporal closures. Incidental fishing mortality by demersal trawlers and auto longliners targeting upper slope teleosts may continue to maintain pressure on the remnant population of Southern Dogfish off southeast Australia.

## Habitat and biology

Southern Dogfish inhabits the upper to mid-continental slope, mainly at depths between 350 and 745 m but with an overall depth range of 215–1400 m. Maximum size is 112 cm total length (TL), with males mature at 82–91 cm TL and females at 87–89 cm TL (Ebert et al. 2021). It has a very low fecundity of one to two pups every two (or possibly three) years and an estimated age at first maturity of >23 years (Whitely 2004).

Longevity and maximum size	Longevity: unknown Max size: 112 cm TL
Age and/or size at maturity (50%)	Males: 82–91 cm TL Females: estimated >23 years, 87–89 cm TL

**CAAB Code:** 37 020011

**Link to IUCN Page:** <https://www.iucnredlist.org/species/41745/124416090>

**Link to page at Shark References:** <http://www.shark-references.com/species/view/Centrophorus-uyato>

## References

- AFMA 2006. *Response to Ministerial Direction- SESSF*. Australian Fisheries Management Authority. Australian Government.
- AFMA. 2012. *Upper-Slope Dogfish Management Strategy. AFMA-managed fisheries. October 2012*. Australian Fisheries Management Authority, Canberra.
- Andrew, N.L., Graham, K.J., Hodgson, K.E. and Gordon, G.N.G. 1997. *Changes after twenty years in relative abundance and size composition of commercial fishes caught during fishery independent surveys on SEF trawl grounds*. NSW Fisheries Final Report Series No. 1 FRDC Project No. 96/139
- Daley, R., Stevens, J.D. and Graham, K. 2002. *Catch analysis and productivity of the deepwater dogfish resource in southern Australia*. Report by CSIRO Marine Research and NSW Fisheries to the Fisheries Research and Development Corporation.

- Duffy, C.A.J. 2007. First record of *Centrophorus harrissoni* from New Zealand, with observations on squamation in Centrophoridae (Squaliformes). *New Zealand Journal of Marine and Freshwater Research* 41: 163–173.
- Graham, K.J., Andrew, N.L. and Hodgson, K.E. 2001. Changes in the relative abundances of sharks and rays on Australian South East Fishery trawl grounds after twenty years of fishing. *Marine and Freshwater Research* 52: 549–561.
- Kyne, P.M., Heupel, M.R., White, W.T. and Simpfendorfer, C.A. 2021. *The Action Plan for Australian Sharks and Rays 2021*. National Environmental Science Program, Marine Biodiversity Hub, Hobart
- Last, P.R. and Stevens, J.D. 2009. *Sharks and Rays of Australia*. Second Edition. CSIRO Publishing, Collingwood.
- Stobutzki, I., Ward, P., Vieira, S. Moore, A., Sahlqvist, P., Leatherbarrow, A., Patterson, H., Barnes, B., Noriega, R. and Rodgers, M. 2011. *Commonwealth trawl and scalefish hook sectors*. In: J. Woodhams, I. Stobutzki, S. Vieira, R. Curtotti and G.A. Begg (eds), Fishery status reports 2010: status of fish stocks and fisheries managed by the Australian Government, pp. 125–192. Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra.
- White, W.T., Guallart, J., Ebert, D.A., Naylor, G.J.P., Verissimo, A., Cottom, C.F., Harris, M., Serena, F and Iglesias, S.P. 2022. Revision of the genus *Centrophorus* (Squaliformes: Centrophoridae): Part 3 – Redescription of *Centrophorus uyato* (Rafinesque) with a discussion of its complicated nomenclatural history. *Zootaxa* 5155(1):001–051.
- Whitely, R. 2004. *Using dorsal spines to age the Australian dogfish Centrophorus harrissoni and Centrophorus uyato*. University of Bangor, Bangor, UK.
- Wilson, D.T., Patterson, H.M., Summerson, R. and Hobsbawn, P.I. 2009. *Information to support management options for upper-slope gulper sharks (including Harrison's dogfish and southern dogfish)*. Final Report to the Fisheries Research and Development Corporation Project No. 2008/65. Bureau of Rural Sciences, Canberra.