


## Gulf Wobbegong, *Orectolobus halei*

<b>Report Card assessment</b>	<b>Sustainable</b>		
IUCN Red List Australian Assessment	Least Concern (Endemic to Australia)	IUCN Red List Global Assessment	Least Concern
Assessors	Huveneers, C., Pollard, D.A., Gordon, I., Flaherty, A.A. & Pogonoski, J.		
Australian Assessors	Kyne, P.M., Heupel, M.R., White, W.T. & Simpfendorfer, C.A. (Shark Action Plan)		
Report Card Remarks	Catch is managed in NSW and low in other states.		

### Summary

The Gulf Wobbegong is endemic to southern Australia. It is a site attached species that occurs at the surface to 195 m depth. It is caught in commercial and recreational fisheries as a target species and as bycatch. In New South Wales (NSW), wobbegong catch declined by more than 50% between 1997–1998 and 2007–2008, after which it stabilised.



This led to all three species of wobbegong occurring in NSW, including the Banded Wobbegong, to be listed as Vulnerable in that state. New NSW management regulations in May 2008 introduced a daily limit of six wobbegongs. As a result, wobbegongs are no longer targeted to the same extent as they used to be in NSW. In other Australian states, wobbegongs are not targeted and catches are low. An IUCN Green List assessment carried out in 2021 found that they were fully recovered from these declines (Huveneers 2021). Therefore, the species is assessed as Least Concern (IUCN) (Kynes et al. 2021) and Sustainable (SAFS).

### Distribution

The Gulf Wobbegong is endemic to Australia. It is found from Norwegian Bay (Western Australia) south through South Australia, Victoria, NSW, and to Hervey Bay (Queensland) (Last and Stevens 2009).

### Stock structure and status

There is currently no information on population size, structure, or trend for the species, but based on catch data from various Australian fisheries there is no reason to believe the population is declining. An IUCN Green List assessment carried out in 2021 found that they were fully recovered from these declines (Huveneers 2021).

## Fisheries

Wobbegongs are caught in trawls, beach seines, gillnets, lobster pots and traps, by hook-and-line, and also by spearfishing. Observed site fidelity in wobbegongs is likely to increase their susceptibility to fishing pressure. Post-release survival is thought to be high. In NSW, three wobbegong species (Spotted Wobbegong, Ornate Wobbegong, and Gulf Wobbegong) are taken in the Ocean Trap and Line Fishery (OTL), in the fish and prawn sectors of the Ocean Trawl Fishery, and in the Estuary General Fishery. The majority of commercial wobbegong catches occur in the OTL Fishery. The NSW total catch of wobbegongs declined from about 120 tonnes in 1990–1991 to about seven tonnes in 2008–2009, largely as a result of reductions in effort. Catches have now stabilized at approximately 20 tonnes per year. Based on species-specific reporting from July 2009, catches of the Gulf Wobbegong ranged from 8.9 to 12.3 tonnes per year, showing no declines. There is also a daily trip limit on wobbegongs and minimum size limit of 180 cm total length. In Queensland, wobbegongs are not targeted but have been recorded in low numbers as bycatch of prawn trawl fisheries (Kyne et al. 2002). Small wobbegongs are sometimes caught by crab pots in Southeast Queensland and Moreton Bay, but are usually discarded (J. Stead, pers. comm., 2007). In South Australia, commercial catches of wobbegong are small ranging from 0.5 to 4 tonnes (A. Tsolos, pers. comm., March 2015). In Western Australian, the Gulf Wobbegong is a small component of the bycatch of the temperate demersal gillnet and demersal longline fisheries. These fisheries catch on average 40 tonnes of wobbegong species per year, and the catch does not show any sign of decline (Department of Fisheries WA Fishery Status Report 1998–99 to 2013–14, Braccini et al. 2014). The Gulf Wobbegong is believed to be a major component of this aggregated catch. However, as all sharks and rays are now commercially protected throughout Western Australia, wobbegongs cannot generally be retained by State-managed commercial fishing vessels unless they are operating in the managed shark fishery.

## Habitat and ecology

The Gulf Wobbegong occurs inshore on the continental shelf to at least 195 m depth (J. Chidlow, pers. comm., 2011). It is found in bays, on macroalgal rocky reef areas, coral reefs, and around offshore islands (Compagno 2001). A survey of wobbegongs showed evidence of site attachment with divers observing individual sharks in exactly the same positions over consecutive dives (Ecology Lab 1991). Maximum size is at least 206 cm total length (TL), with reports to 290 cm TL individuals requiring confirmation (Last and Stevens 2009).

Longevity and maximum size	Longevity: unknown Max size: at least 206 cm TL
Age and/or size at maturity (50%)	Males: 168 cm TL Females: 161 cm TL

**CAAB Code:** 37 013020

**Link to IUCN Page:** <https://www.iucnredlist.org/species/161709/68638176>

**Link to page at Shark References:** <http://shark-references.com/species/view/Orectolobus-halei>

## References

- Braccini, M., McAuley, R. and O'Malley, J. 2014. *Temperate Demersal Gillnet and Demersal Longline Fisheries Status Report*. In: Fletcher, W.J. and Santoro, K. (eds), *Status Reports of the Fisheries and Aquatic Resources of Western Australia 2013/14: The State of the Fisheries*, Department of Fisheries, Western Australia.
- Compagno, L.J.V. 2001. *Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Volume 2. Bullhead, Mackerel and Carpet Sharks (Heterodontiformes, Lamniformes and Orectolobiformes)*. FAO, Rome.

- Ecology Lab. 1991. *A preliminary survey of grey nurse sharks (Carcharias taurus) and wobbegong sharks (Orectolobus spp.) at Seal Rocks, NSW*. The Ecology Lab Pty. Ltd., Balgowlah, New South Wales, Australia.
- Kyne, P.M., Courtney, A., Campbell, M., Chilcott, K., Gaddes, S., Turnbull, C., Van Der Geest, C. and Bennett, M.B. 2002. *An Overview of the Elasmobranch By-catch of the Queensland East Coast Trawl Fishery (Australia)*. Scientific Council Meeting – September 2002. Northwest Atlantic Fisheries Organization Serial No. N4718 NAFO SCR Doc. 02/97.
- Kyne, P.M., Heupel, M.R., White, W.T. and Simpfendorfer, C.A. 2021. *The Action Plan for Australian Sharks and Rays 2021*. National Environmental Science Program, Marine Biodiversity Hub, Hobart
- Last, P.R. and Stevens, J.D. 2009. *Sharks and Rays of Australia*. Second Edition. CSIRO Publishing, Collingwood, Australia.
- Pease, B.C. and Grinberg, A. 1995. *New South Wales Commercial Fisheries Statistics 1940 to 1992*. NSW Fisheries, Cronulla, Australia.